

ASCA Discovery of an X-ray Pulsar in the Error Box of SGR1900+14

K. Hurley, P. Li

University of California, Berkeley, Space Sciences Laboratory, Berkeley, CA 94720-7450

C. Kouveliotou

Universities Space Research Association at NASA Marshall Space Flight Center, ES-84,
Huntsville AL 35812

T. Murakami, M. Ando

ISAS, 3-1-1 Yoshinodai, Sagamihara, Kanagawa, Japan 229

T. Strohmayer

NASA Goddard Space Flight Center, Greenbelt, MD 20771

J. van Paradijs¹, University of Alabama in Huntsville, AL 35899

F. Vrba, C. Luginbuhl

U.S.Naval Observatory, Box 1149, Flagstaff Station, Flagstaff, AZ 86002-1149

A. Yoshida

RIKEN, 2-1, Hirowawa, Wako, Saitama, 351-01, Japan

I. Smith

Department of Space Physics and Astronomy, Rice University, MS-108, 6100 South Main,
Houston, TX 77005-1892

S. Kulkarni

California Institute of Technology, Department of Astronomy, 105-24, Pasadena, CA 91125

ABSTRACT

We present a 2 - 10 keV ASCA observation of the field around the soft gamma repeater SGR1900+14. One quiescent X-ray source was detected in this observation, and it was in the SGR error box. In 2 - 10 keV X-rays, its spectrum may be fit by a power law with index -2.2, and its unabsorbed flux is $9.6 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$. We also find a clear 5.16 s period. The properties of the three well-studied soft gamma repeaters are remarkably similar to one another, and provide evidence that all of them are associated with young, strongly magnetized neutron stars in supernova remnants.

Subject headings: gamma rays: bursts — stars: neutron — X-rays: stars — supernova remnants

¹Astronomical Institute ‘Anton Pannekoek’, University of Amsterdam, The Netherlands

1. Introduction

Four soft gamma repeaters (SGRs) are now known to exist with certainty, and there is good evidence that all of them are associated with supernova remnants (SNRs). The position of SGR0525-66 is consistent with that of the N49 supernova remnant in the Large Magellanic Cloud (Cline et al. 1982). SGR1806-20 (Atteia et al. 1987) is associated with the SNR G10.0-0.3, located towards the galactic center (Kulkarni & Frail 1993; Kouveliotou et al. 1994; Kulkarni et al. 1994; Murakami et al. 1994). SGR1627-41 may be associated with G337.0-0.1 (Woods et al., 1998; Hurley et al. 1998a,b). SGR1900+14 lies close to G42.8+0.6 (Kouveliotou et al. 1994; Hurley et al. 1998c). Three of the four are quiescent soft X-ray point sources: SGR0525-66 (Rothschild, Kulkarni, & Lingener 1994), SGR1806-20 (Murakami et al. 1994), and SGR1900+14 (Hurley et al. 1994, 1998c). Finally two of the SGRs, 0525-66 (Mazets et al. 1979, Barat et al. 1979) and 1900+14 (Hurley et al. 1998d) display periodicities in their bursting emission, and one, 1806-20 (Kouveliotou et al. 1998a) in its quiescent emission, indicating that they are neutron stars.

In this paper we present soft X-ray observations of the quiescent X-ray source associated with SGR1900+14 by the Advanced Satellite for Cosmology and Astrophysics (ASCA) which indicate that it, too displays a periodicity in its quiescent emission. Furthermore, its X-ray spectrum is similar to that of SGR1806-20. In another paper (Hurley et al. 1998c) we present evidence, based on burst localization by the interplanetary network, that this X-ray source is indeed associated with the SGR. Thus SGR1900+14 is similar to the other SGRs in many respects, pointing towards a unified model for these sources.

Kouveliotou et al. (1994) suggested that SGR1900+14 might be associated with one of two SNRs, G43.9+1.6 or G42.8+0.6. Both the ROSAT sky survey (Vasisht et al. 1994) and a pointed observation at the position of G42.8+0.6 (Hurley et al. 1996) indicated that a point-like quiescent X-ray source was present near this SNR. This source was also associated with one of the two network synthesis error boxes for this SGR (Hurley et al. 1994). No ROSAT source was detected in the second network synthesis error box (Li et al. 1997), and the more recent localization for SGR1900+14 (Hurley et al. 1998c) indicates that the first error box is indeed the one which contains the SGR and the quiescent X-ray source. Optical and infrared observations of the X-ray position revealed a peculiar double M star system within the X-ray source error box (Vrba et al. 1996). A rough estimate of the probability of finding a random X-ray source of any intensity in the error box for this SGR may be obtained from the statistics of the WGA catalog (White, Giommi, and Angelini 1995). There are $\sim 10,000$ sources located at galactic latitudes $|b| < 20^\circ$, or 1.12×10^{-3} sources/ sq. arcmin., and the error box area is ~ 1.6 sq. arcmin. (Hurley et al. 1998c). Thus the probability is ~ 0.0018 . This estimate makes no correction for the sky exposure as a function of galactic latitude, and encompasses the widely varying sensitivities of the ROSAT pointed observations in the public domain (typical observation times 5000 - 30,000 s). However, as we show below, due to the fact that the period of the quiescent source is identical to that found in the giant flare of 1998 August 27 (Hurley et al. 1998d), the association between the SGR and the X-ray source does not depend strongly on this probability.

2. ASCA Observations

The position of the quiescent X-ray point source associated with SGR1900+14, $\alpha(2000) = 19^{\text{h}}07^{\text{m}}14^{\text{s}}$, $\delta(2000) = 9^{\circ}19'19''$, was known from previous ROSAT observations (Hurley et al. 1994). ASCA was pointed to within $\sim 0.1^{\circ}$ of this position on 1998 April 30. (An earlier attempt at an observation on 1997 October 22 was incorrectly pointed). The observation began around 20:55 UT and lasted 2.3 days, resulting in 74 ksec of on-source exposure with the Solid State Imaging Spectrometer (SIS), and 84.6 ksec with the Gas Imaging Spectrometer (GIS). We used the standard screening criteria for such parameters as Earth elevation angle, South Atlantic Anomaly, and cutoff rigidity to extract photons. These are described in the ASCA Data Reduction Guide, Version 2². No bursts from the source were observed by *Ulysses*, BATSE, KONUS, or ASCA during the observation. One quiescent source was detected, at $\alpha(2000) = 19^{\text{h}}07^{\text{m}}16.4^{\text{s}}$, $\delta(2000) = +09^{\circ}19'44.1''$ with an error radius 40'', consistent with the position of the ROSAT source. The source location was determined with the Ximage source detection tool, which uses a sliding window technique.

The region used for spectral analysis consisted of a 6' radius circle centered at the source position; background was taken from the same observation, using a 6' radius circle at a region where no source was present, as determined by the same sliding window technique. Spectral fitting was done using XSPEC and three trial functions: blackbody, thermal bremsstrahlung, and a power law, all with absorption. Using GIS2 and GIS3 data, a power law gave the best fit, with a reduced χ^2 of 1.48 for 182 degrees of freedom. The best reduced χ^2 for the blackbody fit was 4.38, and that for the bremsstrahlung was 2.04. We therefore adopt the power law fit; the best fit photon power law index was 2.25 ± 0.04 , and the absorption was $n_{\text{H}} = (2.16 \pm 0.07) \times 10^{22} \text{ cm}^{-2}$. This corresponds to an unabsorbed 2 - 10 keV flux of $(9.6 \pm 0.7) \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$. The fit is shown in figure 1.

Previously, the distance to SGR1900+14 had been estimated as 5 kpc using the Σ -D relation (e.g. Vasisht et al. 1994). The measurement of n_{H} now permits an independent estimate. Using the W3nH tool available on-line from the HEASARC³ we obtain ≈ 5.7 kpc. Both methods are subject to uncertainties; in what follows, we adopt a distance of 5 kpc.

To search for periodicity, light curves were constructed with 0.5 s binning from the sum of the GIS2 and GIS3 data, by extracting $\approx 1 - 10$ keV counts from a 4' radius circular region around the source. The power spectrum is shown in figure 2. A significant barycenter-corrected period was found at 5.1589715 ± 0.0000008 s. The distribution of powers closely follows a χ^2 distribution with two degrees of freedom. Taking into account the number of frequencies searched, the chance probability of finding a peak this strong or stronger in the power spectrum may be estimated at 4.7×10^{-14} , a conservative number which does not take the presence of the harmonics into account.

²available on-line at <http://heasarc.gsfc.nasa.gov/docs/asca/abc/abc.html>

³http://heasarc.gsfc.nasa.gov/docs/frames/mb_w3browse.html

A folded light curve is shown in figure 3. This period agrees well with that found for the SGR in outburst on 1998 August 27 (Hurley et al. 1998d), making the identification of the quiescent source and the SGR quite secure.

3. Discussion and Conclusion

The ASCA observations of SGR1900+14 may be used to estimate the magnetic field of the neutron star in two ways. First, we can use the association between the quiescent X-ray source and G42.8+0.6. The observable lifetime of SNRs is less than $\sim 20,000$ years (Braun, Goss, and Lyne 1989). The characteristic age τ of a rotating, magnetized neutron star is related to the period P and its derivative $\dot{P}$ by $\tau = P/2\dot{P}$, while P and $\dot{P}$ are related to the field strength B by $P\dot{P} = AB^2$, where $A = 9.8 \times 10^{-40} \text{ s G}^{-2}$ (e.g. Lyne, Manchester, and Taylor 1985). Thus $B = \sqrt{P^2/2A\tau}$. For SGR1900+14, we obtain $B = 2 \times 10^{14} \text{ G}$ for an age of 10^4 y , remarkably similar to estimates made from the energy contained in the soft tail of the giant flare of 1998 August 27 (Hurley et al. 1998d) and from direct measurement of $\dot{P}$ (Kouveliotou et al. 1998b). It is also comparable to the field strength of SGR1806-20, estimated from direct measurements of P and $\dot{P}$ for that source (Kouveliotou et al. 1998a). This suggests that SGR behavior can be described by the magnetar model (Thompson and Duncan 1995). If this is the case, a second estimate of B comes from one of the predictions of that model, namely that persistent soft X-ray emission is powered by crustal magnetic energy: $\frac{B_{crust}^2}{8\pi} 4\pi R_*^2 \Delta R \gtrsim L_x \tau$, where R_* is the radius of the neutron star (10 km), ΔR is the thickness of the crust ($\sim 1 \text{ km}$), and we take $L_x = 2.7 \times 10^{34} \text{ erg s}^{-1}$ for a distance of 5 kpc and $\tau = 10,000 \text{ y}$. We obtain $B_{crust} \gtrsim 4 \times 10^{14} \text{ G}$. The two estimates would agree ($B \sim 2.5 \times 10^{14} \text{ G}$) for an age $\tau \sim 7100 \text{ y}$, or about 70% of the lifetime of a typical magnetar, in the model.

Table 1 compares the properties of the three SGRs and their soft X-ray counterparts. The X-ray spectrum of SGR0525 is unknown. However, the spectra of SGR1806 and SGR1900 have identical power law indices; their 2 - 10 keV luminosities agree to within an order of magnitude when their estimated distances are taken into account. The periods of all three sources agree to within a factor of ~ 1.6 . The four known SGRs therefore share many similarities – in their periods, the properties of their quiescent soft X-ray sources, their ages, their magnetic fields, and their associations with supernova remnants. This provides further evidence that, as a class, the SGRs are strongly magnetized young neutron stars whose behavior can be described by the magnetar model of Thompson and Duncan (1995).

KH and PL are grateful to NASA for support under the ASCA AO-6 Guest Investigator Program. CK acknowledges support under the CGRO guest investigator program (NAG 5-2560). We are also grateful to Dr. K. Mitsuda who kindly provided his computer code to confirm the period.

REFERENCES

- Atteia, J.-L. et al. 1987, ApJ 320, L105
- Barat, C. et al. 1979, A&A 79, L24
- Braun, R., Goss, W., and Lyne, A. 1983, ApJ 340, 355
- Cline, T. et al. 1982, ApJ 255, L45
- Hurley, K. et al. 1994, ApJ 431, L31
- Hurley, K. et al. 1996, ApJ 463, L13
- Hurley, K., Kouveliotou, C., Kippen, R.M., and Woods, P. 1998a, IAU Circ. 6948
- Hurley, K., Cline, T., Butterworth, P., Mazets, E., and Golenetskii, S. 1998b, GCN GRB Observation Report 122
- Hurley, K. et al. 1998c, ApJ companion paper
- Hurley, K. et al. 1998d, Nature accepted
- Kouveliotou, C. et al. 1994, Nature 328, 125
- Kouveliotou, C. et al. 1998a, Nature 393, 235
- Kouveliotou, C. et al. 1998b, ApJ companion paper
- Kulkarni S. and Frail, D. 1993, Nature 365, 33
- Kulkarni S. et al. 1994, Nature 368, 129
- Li, P., Hurley, K., Vrba, F., Kouveliotou, C., Meegan, C., Fishman, G., Kulkarni, S., and Frail, D. 1997, ApJ 490, 823
- Lyne, A., Manchester, R., and Taylor, J. 1985, MNRAS 213, 613
- Mazets, E. et al., 1979, Nature 282, 587
- Murakami, T. et al. 1994, Nature 368, 127
- Murakami, T. 1995, Ap&SS 231, 57
- Murakami, T. et al. 1996, in Gamma-Ray Bursts, 3rd Huntsville Symposium, Eds. C. Kouveliotou, M. Briggs, and G. Fishman, AIP Conf. Proc. 384 (AIP: New York), p. 961
- Rothschild, R., Kulkarni, S., and Lingenfelter, R. 1994, Nature 368, 432
- Thompson, C., and Duncan, R. 1995, MNRAS 275, 255
- Vasisht, G. et al. 1994, ApJ 431, L35
- Vrba, F. et al. 1996, ApJ 468, 225
- White, N., Giommi, P., and Angelini, L., The WGA Catalog of ROSAT Point Sources, available on-line through the HEASARC (<http://lheawww.gsfc.nasa.gov/users/white/wgacat/wgacat.html>)

Woods, P., Kippen, R.M., vanParadijs, J., Kouveliotou, C., McCollough, M., and Hurley, K. 1998, IAU Circ. 6948




Fig. 1.— Best fit power law spectrum and residuals for the GIS2 and GIS3 data. The large residuals around 1 keV are believed to be caused by a small gain shift.




Fig. 2.— Power spectrum of the 2 - 10 keV soft X-ray source associated with SGR1900+14. The 0.19384-Hz pulsed signal and its first and second harmonics are well above the noise level.




Fig. 3.— Epoch folded pulse profile of SGR1900+14 from the 2 - 10 keV ASCA data.

Table 1. *Properties of three SGRs*

SGR	Soft X-ray spectrum	Flux, erg cm ⁻² s ⁻¹	Period, s	Distance, kpc
0525-66 ^{1,2,3,4}	? ⁵	2x10 ⁻¹²	8	55
1806-20 ^{2,3,6}	E ^{-2.2}	1x10 ⁻¹¹	7.5	~ 15(?)
1900+14	E ^{-2.2}	9.6x10 ⁻¹²	5.2	~ 5(?)

¹Rothschild, Kulkarni, and Lingenfelter 1994

²Murakami 1995

³Murakami et al. 1996

⁴Barat et al. 1979

⁵Source was detected by the ROSAT HRI, but not resolved by ASCA; no spectral measurement available. Flux is for 0.1 - 2.4 keV

⁶Kouveliotou et al. 1998a